


January 2015 Volume 64 Issue 1

Gut

An International Journal of
Gastroenterology
and Hepatology

gut.bmj.com

BMJ



Caption: Winter wonderland but very hot January issue of *Gut*. Every article in the January issue is freely available.

An international journal of gastroenterology and hepatology

Impact Factor: 13.319

Editor

Emad El-Omar (UK)

Deputy Editor (Hepatology)

Alexander Gerbes (Germany)

Deputy Editor (Luminal

Gastroenterology)

William Grady (USA)

Deputy Editor (Endoscopy)

Thomas Rösch (Germany)

Associate Editors

Matias Avila (Spain)

Martin Blaser (USA)

Guy Boeckxstaens (Belgium)

Thomas Gress (Germany)

Arthur Kaser (UK)

Ernst Kuipers (The Netherlands)

Jake Liang (USA)

Julian Panes (Spain)

Herbert Tilg (Austria)

Fabien Zoulim (France)

Editorial Office

Gut

BMJ Journals

BMA House, Tavistock Square

London, WC1H 9JR, UK

T: +44 (0)20 7383 6394

F: +44 (0)20 7383 6668

E: info.gut@bmj.com

Twitter: @Gut_BMJ

ISSN: 0017-5749 (print)

ISSN: 1468-3288 (online)

Disclaimer: *Gut* is owned and published by BMJ Publishing Group Ltd, a wholly owned subsidiary of the British Medical Association and the British Society of Gastroenterology. The owners grant editorial freedom to the Editor of *Gut*. *Gut* follows guidelines on editorial independence produced by the World Association of Medical Editors and the code on good publication practice of the Committee on Publication Ethics.

Gut is intended for medical professionals and is provided without warranty, express or implied. Statements in the Journal are the responsibility of their authors and advertisers and not authors' institutions, the BMJ Publishing Group, the British Society of Gastroenterology or the BMA unless otherwise specified or determined by law. Acceptance of advertising does not imply endorsement.

To the fullest extent permitted by law, the BMJ Publishing Group shall not be liable for any loss, injury or damage resulting from the use of *Gut* or any information in it whether based on contract, tort or otherwise. Readers are advised to verify any information they choose to rely on.

Copyright © 2014 BMJ Publishing Group Ltd and British Society of Gastroenterology. All rights reserved; no part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise without prior permission.

Gut is published by BMJ Publishing Group Ltd, typeset by Techset and printed in the UK on acid-free paper.

Gut (ISSN: 0017-5749) is published monthly by BMJ Publishing Group and is distributed in the USA by Air Business Ltd. Periodicals postage paid at Jamaica NY 11431 POSTMASTER: send address changes to *Gut*, Air Business Ltd, c/o Worldnet Shipping Inc., 156-15, 146th Avenue, 2nd Floor, Jamaica, NY 11434, USA.

Contents

Volume 64 Issue 1 | **GUT** January 2015

Leading article

- 1 Novel strategies for liver therapy using stem cells

T Rashid, T Takebe, H Nakauchi

Commentaries

- 5 Does a prior diagnosis of Barrett's oesophagus influence risk of dying from oesophageal adenocarcinoma?

D C Whiteman

- 6 Treatment of paediatric cholestasis due to canalicular transport defects: yet another step forward

J J G Marin, R H J Houwen

- 8 Targeting of gut specific leucocyte recruitment in IBD by vedolizumab

B Eksteen

Oesophagus

- 11 The human squamous oesophagus has widespread capacity for clonal expansion from cells at diverse stages of differentiation

OPEN ACCESS

M Barbera, M di Pietro, E Walker, C Brierley, S MacRae, B D Simons, P H Jones, J Stingl, R C Fitzgerald

- 20 Oesophageal adenocarcinoma and prior diagnosis of Barrett's oesophagus: a population-based study

S K Bhat, D T McManus, H G Coleman, B T Johnston, C R Cardwell, Ú McMenamin, F Bannon, B Hicks, G Kennedy, A T Gavin, L J Murray

- 26 TBL1XR1 promotes lymphangiogenesis and lymphatic metastasis in esophageal squamous cell carcinoma

L Liu, C Lin, W Liang, S Wu, A Liu, J Wu, X Zhang, P Ren, M Li, L Song

Stomach

- 37 OCT1 is a determinant of synbindin-related ERK signalling with independent prognostic significance in gastric cancer

OPEN ACCESS

J Qian, X Kong, N Deng, P Tan, H Chen, J Wang, Z Li, Y Hu, W Zou, J Xu, J-Y Fang

Endoscopy

- 49 The combination of autofluorescence endoscopy and molecular biomarkers is a novel diagnostic tool for dysplasia in Barrett's oesophagus

OPEN ACCESS

M di Pietro, D F Boerwinkel, M K Shariff, X Liu, E Telakis, P Lao-Sirieix, E Walker, G Couch, L Mills, T Nuckcheddy-Grant, S Slininger, M O'Donovan, M Visser, S L Meijer, P V Kaye, L Wernisch, K Ragnanath, J J G H M Bergman, R C Fitzgerald

- 57 Long-term adenoma recurrence following wide-field endoscopic mucosal resection (WF-EMR) for advanced colonic mucosal neoplasia is infrequent: results and risk factors in 1000 cases from the Australian Colonic EMR (ACE) study

A Moss, S J Williams, L F Hourigan, G Brown, W Tam, R Singh, S Zanati, N G Burgess, R Sonson, K Byth, M J Bourke

Inflammatory bowel disease

- 66 XIAP variants in male Crohn's disease



Y Zeissig, B-S Petersen, S Milutinovic, E Bosse, G Mayr, K Peuker, J Hartwig, A Keller, M Kohl, M W Laass, S Billmann-Born, H Brandau, A C Feller, C Röcken, M Schrappe, P Rosenstiel, J C Reed, S Schreiber, A Franke, S Zeissig

- 77 Vedolizumab affects antibody responses to immunisation selectively in the gastrointestinal tract: randomised controlled trial results

T Wyant, T Leach, S Sankoh, Y Wang, J Paolino, M F Pasetti, B G Feagan, A Parikh

Neurogastroenterology

- 84 Increased colonic bile acid exposure: a relevant factor for symptoms and treatment in IBS

A Bajor, H Törnblom, M Rudling, K-A Ung, M Simrén

Gut microbiota

- 93 Diets that differ in their FODMAP content alter the colonic luminal microenvironment

E P Halmos, C T Christophersen, A R Bird, S J Shepherd, P R Gibson, J G Muir

MORE CONTENTS ►



This article has been chosen by the Editor to be of special interest or importance and is freely available online.



This article has been made freely available online under the BMJ Journals Open Access scheme.

See <http://gut.bmj.com/site/about/guidelines.xhtml#open>



COMMITTEE ON PUBLICATION ETHICS

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics

<http://publicationethics.org/>



When you have finished with this magazine please recycle it.

Colon

- 101** Role of tumour molecular and pathology features to estimate colorectal cancer risk for first-degree relatives

A K Win, D D Buchanan, C Rosty, R J MacInnis, J G Dowty, G S Dite, G G Giles, M C Southey, J P Young, M Clendenning, M D Walsh, R J Walters, A Boussioutas, T C Smyrk, S N Thibodeau, J A Baron, J D Potter, P A Newcomb, L L Marchand, R W Haile, S Gallinger, N M Lindor, J L Hopper, D J Ahnen, M A Jenkins

- 111** A candidate gene study of capecitabine-related toxicity in colorectal cancer identifies new toxicity variants at *DPYD* and a putative role for *ENOSF1* rather than *TYMS*



D Rosmarin, C Palles, A Pagnamenta, K Kaur, G Pita, M Martin, E Domingo, A Jones, K Howarth, L Freeman-Mills, E Johnstone, H Wang, S Love, C Scudder, P Julier, C Fernandez-Rozadilla, C Ruiz-Ponte, A Carracedo, S Castellvi-Bel, A Castells, A Gonzalez-Neira, J Taylor, R Kerr, D Kerr, I Tomlinson

- 121** An updated Asia Pacific Consensus Recommendations on colorectal cancer screening

J J Y Sung, S C Ng, F K L Chan, H M Chiu, H S Kim, T Matsuda, S S M Ng, J Y W Lau, S Zheng, S Adler, N Reddy, K G Yeoh, K K F Tsoi, J Y L Ching, E J Kuipers, L Rabeneck, G P Young, R J Steele, D Lieberman, K L Goh

Pancreas

- 133** Simvastatin is associated with reduced risk of acute pancreatitis: findings from a regional integrated healthcare system

B U Wu, S J Pandol, I-L A Liu

Hepatology

- 139** Anti-HBV DNA vaccination does not prevent relapse after discontinuation of analogues in the treatment of chronic hepatitis B: a randomised trial—ANRS HB02 VAC-ADN

H Fontaine, S Kahi, C Chazallon, M Bourguine, A Varaut, C Buffet, O Godon, J F Meritet, Y Saïdi, M L Michel, D Scott-Algara, J P Aboulker, S Pol, for the ANRS HB02 study group

- 147** Functional analysis of *ABCB4* mutations relates clinical outcomes of progressive familial intrahepatic cholestasis type 3 to the degree of MDR3 floppase activity

R Gordo-Gilart, S Andueza, L Hierro, P Martínez-Fernández, D D'Agostino, P Jara, L Alvarez

- 156** Epigenetic modification of MiR-429 promotes liver tumour-initiating cell properties by targeting Rb binding protein 4

L Li, J Tang, B Zhang, W Yang, M Liu Gao, R Wang, Y Tan, J Fan, Y Chang, J Fu, F Jiang, C Chen, Y Yang, J Gu, D Wu, L Guo, D Cao, H Li, G Cao, M Wu, Michael Q Zhang, L Chen, H Wang

- 168** Therapeutic targeting of GSK3 β enhances the Nrf2 antioxidant response and confers hepatic cytoprotection in hepatitis C

Y Jiang, H Bao, Y Ge, W Tang, D Cheng, K Luo, G Gong, R Gong

Editor's quiz: GI snapshot

- 180** Endoscopic submucosal dissection of an unusual flat rectal neoplasm

F Azzolini, P Cecinato, V Iori, L De Marco, R Zecchini, C Fodero, C Tioli, R Sassatelli

JournalScan

- 181** GI highlights from the literature

M H McLean

PostScript

- 184** Letters

Receive regular table of contents by email.
Register using this QR code.





OPEN ACCESS

ORIGINAL ARTICLE

OCT1 is a determinant of synbindin-related ERK signalling with independent prognostic significance in gastric cancer

Jin Qian,¹ Xuan Kong,¹ Niantao Deng,² Patrick Tan,^{2,3} Haoyan Chen,¹ Jilin Wang,¹ Zhaoli Li,⁴ Ye Hu,¹ Weiping Zou,⁵ Jie Xu,¹ Jing-Yuan Fang¹

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/gutjnl-2013-306584>).

¹State Key Laboratory of Oncogenes and Related Genes, Division of Gastroenterology and Hepatology, Renji Hospital, School of Medicine, Shanghai Jiao Tong University, Shanghai Cancer Institute, Shanghai Institute of Digestive Disease, Shanghai, China

²Cancer and Stem Cell Biology Program, Duke-NUS Graduate Medical School, Singapore, Singapore

³Cancer Therapeutics and Stratified Oncology, Genome Institute of Singapore, Singapore, Singapore

⁴Harbin Veterinary Research Institute, Chinese Academy of Agricultural Sciences, Harbin, China

⁵Department of Surgery, University of Michigan, Ann Arbor, Michigan, USA

Correspondence to

Dr Jing-Yuan Fang;
jingyuanfang@yahoo.com
and Dr Jie Xu;
jiexu@yahoo.com

Received 11 December 2013

Accepted 17 March 2014

Published Online First

9 April 2014



Open Access
Scan to access more
free content



CrossMark

To cite: Qian J, Kong X, Deng N, et al. *Gut* 2015;64:37–48.

ABSTRACT

Objective Octamer transcription factor 1 (OCT1) was found to be expressed in intestinal metaplasia and gastric cancer (GC), but the exact roles of OCT1 in GC remain unclear. The objective of this study was to determine the functional and prognostic implications of OCT1 in GC.

Design Expression of OCT1 was examined in paired normal and cancerous gastric tissues and the prognostic significance of OCT1 was analysed by univariate and multivariate survival analyses. The functions of OCT1 on synbindin expression and extracellular signal-regulated kinase (ERK) phosphorylation were studied in vitro and in xenograft mouse models.

Results The OCT1 gene is recurrently amplified and upregulated in GC. OCT1 overexpression and amplification are associated with poor survival in patients with GC and the prognostic significance was confirmed by independent patient cohorts. Combining OCT1 overexpression with American Joint Committee on Cancer staging improved the prediction of survival in patients with GC. High expression of OCT1 associates with activation of the ERK mitogen-activated protein kinase signalling pathway in GC tissues. OCT1 functions by transactivating synbindin, which binds to ERK DEF domain and facilitates ERK phosphorylation by MEK. OCT1-synbindin signalling results in the activation of ERK substrates ELK1 and RSK, leading to increased cell proliferation and invasion. Immunofluorescent study of human GC tissue samples revealed strong association between OCT1 protein level and synbindin expression/ERK phosphorylation. Upregulation of OCT1 in mouse xenograft models induced synbindin expression and ERK activation, leading to accelerated tumour growth in vivo. **Conclusions** OCT1 is a driver of synbindin-mediated ERK signalling and a promising marker for the prognosis and molecular subtyping of GC.

INTRODUCTION

Stomach adenocarcinoma, or gastric cancer (GC), is the fourth most common cancer and the second highest cause of cancer-related mortality worldwide.¹ The prognosis of patients with GC continues to be dismal, despite improving surgical and adjuvant treatment approaches, with a 5-year overall survival less than 25%.² It is of great clinical importance to identify genes that control the severity of GC and present predictive value for prognosis.^{3,4} In a systematic study of molecular signatures in GC, mitogen-activated protein kinase (MAPK)

Significance of this study

What is already known on this subject?

- Octamer transcription factor 1 (OCT1) is a homologue of OCT4 pluripotency factor, and it is expressed in intestinal metaplasia foci and in gastric cancers (GCs).
- The mitogen-activated protein kinase (MAPK) pathway is one of the most frequently altered signalling pathways in GC.
- Synbindin is a spatial regulator of the extracellular signal-regulated kinases (ERK)/MAPK signalling pathway.

What are the new findings?

- The OCT1 gene is recurrently amplified in the genome of GC, and OCT1 amplification displays mutual exclusivity with KRAS and FGFR2 genes in the receptor tyrosine kinase (RTK) pathway.
- OCT1 gene amplification and upregulation are associated with poor survival of patients with GC, and the association is supported by independent datasets.
- OCT1 transactivates synbindin, which binds to the ERK DEF domain and potentiates ERK phosphorylation by MEK1, leading to activation of the ERK substrates ELK1 and RSK. The expression of OCT1 strongly correlates with the levels of synbindin and ERK phosphorylation in GC tissues.
- OCT1 upregulation suppresses apoptosis and enhances proliferation and invasion of GC cells. Knockdown of synbindin expression blocked the pro-malignant effects of OCT1.
- In xenograft tumour models, OCT1 substantially promoted synbindin expression and ERK activation, leading to accelerated tumour growth in vivo.

was found as the most frequently activated pathway in this deadly disease.⁵ In support of this, several components of the receptor tyrosine kinase (RTK)/RAS/MAPK pathway were found to be frequently amplified in GC.^{6,7} Since MAPK signalling is also controlled by spatiotemporal regulatory mechanisms,^{8,9} it is of interest to test if alternative pathways may contribute to MAPK deregulation in GC.

Significance of this study

How might it impact on clinical practice in the foreseeable future?

- Amplification of OCT1 contributes to ERK/MAPK activation, and it may join other components in the RTK pathway to mark patients with GC who are potentially treatable by RTK/RAS directed therapies (up to 46% of the GC population by simplified estimation).
- OCT1 is an independent prognostic factor in GC, and combining OCT1 expression with American Joint Committee on Cancer staging could further improve the prediction of patient survival. OCT1 gene amplification and upregulation can be used as a promising biomarker for the prognosis of GC.

Octamer transcription factor 1 (OCT1) (POU2F1) belongs to the POU homeodomain family of transcription factors.¹⁰ This protein activates or represses the transcription of various genes, such as the immunoglobulin genes in B cells¹¹ and several interleukins.¹² Accumulating data suggest that OCT1 may contribute to the malignant transformation process, and loss of OCT1 inhibits oncogenic transformation in mouse embryonic fibroblasts and tumorigenicity in p53-deficient mice.¹³ Moreover, OCT1 is believed to facilitate the tumorigenesis of pancreatic and intestinal cancer cells through transactivating the CDX2 gene.¹⁴ Interestingly, OCT1 has been reported to be positive in 87% of intestinal metaplasia foci (a preneoplastic lesion) and in 74% of gastric carcinomas in one series.¹⁵ Nonetheless, in GC cells OCT1 was found without the ability to transactivate CDX2, although it could bind to the CDX2 promoter.¹⁵ Despite recent efforts to understand the activities of OCT1 as a transcriptional factor,¹⁶ little is known about the roles of OCT1 in gastric carcinogenesis and it remains unclear whether OCT1 expression may associate with any clinicopathological features of GC.

Here we report OCT1 as an independent prognostic marker for GC and it regulates the extracellular signal-regulated kinase (ERK) MAPK signalling by transactivating synbindin. Synbindin is a core subunit of the trafficking protein particle complex and it is involved in the targeting and fusion of vesicles from endoplasmic reticulum to Golgi apparatus.¹⁷ We previously found that synbindin functions as a scaffold protein that binds to MEK and ERK2, facilitating MEK-dependent phosphorylation and activation of ERK2 on the Golgi.⁹ In this study, we further show that OCT1 binds to and transactivates the promoter of synbindin, thereby inducing synbindin-mediated activation of ERK signalling in GC. We obtained *in vitro* and *in vivo* data supporting the major regulatory effects of OCT1 on ERK signalling and the aggressive behaviours of GC cells. Importantly, we demonstrate that OCT1 amplification and upregulation associate with poor outcome in patients with GC. By these approaches we aim to elucidate the functional and prognostic implications of OCT1 in GC.

MATERIALS AND METHODS**Immunofluorescence**

Tissue specimens were from patients (89 cases for immunofluorescence and 10 for fluorescent *in situ* hybridization (FISH)) who underwent surgery at the Shanghai Renji Hospital from July 2003 to January 2009. The protocol had the approval of the

Ethics Committee of the Shanghai Jiao-Tong University School of Medicine, Renji Hospital, and the research was carried out according to the provisions of the Helsinki Declaration of 1975. Written informed consent was obtained from all participants involved in the study, and clinical information was collected with Institutional Review Board approval. Meanwhile, 89 specimens of adjacent tissues were taken from these patients as the paired controls. The tissue sections were deparaffinised in xylene and rehydrated using a graded series of ethanol. All slides were treated with NaBH₄ to suppress autofluorescence of tissues. The expression levels of OCT1 and synbindin were probed with the primary antibodies (OCT1, dilution 1:40; synbindin, dilution 1:40) according to the manufacturer's instructions. Secondary antibodies (Alexa488-anti-mouse and Alexa546-anti-rabbit) were used to label synbindin and OCT1, respectively. Protein expression was quantified based on staining intensity.

Fluorescent *in situ* hybridisation

The FISH protocol to detect gene copy number variation (CNV) in formalin-fixed paraffin embedded tissues has been described in detail previously.¹⁸ Briefly, the OCT1 DNA probes were labelled using spectrum green and control probes in spectrum orange (centromeric CEP probe for chromosomes 1) (Abbott Molecular Inc, Des Plaines, Illinois, USA). Hybridised slides were counterstained with DAPI and analysed using Zeiss LSM710 confocal microscope (Carl Zeiss, Regensburg, Germany).

Additional materials and methods including chromatin immunoprecipitation (ChIP), cell viability and invasion assays, flow cytometry, gene set enrichment analysis (GSEA), CNV analysis, TF binding identification, Western blotting, and *in vivo* experiments can be found in the online supplementary information.

RESULTS**OCT1 expression and CNV in GC**

We first investigated OCT1 expression level in gastric adenocarcinoma (90 cases) and their adjacent normal controls by immunofluorescence histochemistry. As a result, OCT1 was weakly or moderately detected in normal gastric mucosa (figure 1A), but its expression level was substantially increased in GC cells ($p<0.0001$, figure 1B,C). Notably, OCT1 was strongly positive in the nucleus of GC cells, suggesting its active status as a transcriptional factor (figure 1D). In addition, we detected increased OCT1 expression in gastric tissues with intestinal metaplasia (see online supplementary figure S1A), which confirmed previous findings.¹⁵

Further, we analysed the CNV of OCT1 in GC tissues using data from The Cancer Genome Atlas (TCGA) GC cohort,¹⁹ the Singapore⁶ and the VUMC cohorts.²⁰ The GISTIC 2.0 package was employed to identify focal alteration events based on Affymetrix SNP6 microarray data (TCGA and Singapore datasets), while the CGHcall algorithm was used to analyse aCGH data (VUMC dataset). The OCT1 (POU2F1) gene is located in a recurrently amplified region in chromosome 1 (see online supplementary figure S1B), with 31 of 305 patients (10.16%) showing focal amplification (q value <0.001) but only one patient with deletion (figure 1E). This result was confirmed by the Singapore GC patient cohort, which reported 18 in 193 patients with GC (9.33) with focal amplification of OCT1 gene (figure 1E). In fact, algorithms that identify focal and broad (q value <0.25) CNV events show much higher frequency of OCT1 amplifications, ranging from 22.2% (40 of 180 cases in VUMC cohort) to 32.1% (98 of 305 cases in TCGA cohort) (figure 1E, see online supplementary figure S1C–E). We




Figure 1 Amplification and overexpression of octamer transcription factor 1 (OCT1) gene in gastric cancer (GC). (A) Expression of OCT1 in normal gastric tissues as determined by immunofluorescence. Scale bars indicate 10µm in all panels. (B) Immunostain of OCT1 in gastric cancer tissues. (C) Statistical analysis of OCT1 expression in normal and GC tissues ($P < 0.0001$, Mann Whitney test). (D) The immunofluorescent image shows OCT1 (in green) and cell nucleus (in blue) OCT1 is strongly expressed in the cell nucleus (red arrows). (E) The frequency of OCT1 amplification in three independent GC cohorts as indicated. (F) OCT1 genomic amplification confirmed by fluorescence in-situ hybridization (FISH) assay. Green signals indicate the OCT1 FISH probe, and red signals probe to centromere 1. Scale bars indicate 5µm in all panels. (G) OCT1 mRNA levels were significantly higher in samples with OCT1 gained CNA compared with the samples without CNV in the TCGA dataset ($P < 0.01$, 2-sided t-test). (H) Relationship between the CNVs of OCT1 and frequently altered genes in Receptor tyrosine kinase (RTK) pathway. The mutual exclusivity between the CNVs of OCT1 (POU2F1) and other genes was determined by dimension reduction permutation (DRP) algorithm (* $P < 0.05$).




Figure 2 Octamer transcription factor 1 (OCT1) gene amplification and upregulation are associated with poor survival in patients with GC. (A) Kaplan-Meier survival analysis of GC patients stratified by OCT1 expression level (median split). (B) Kaplan-Meier survival analysis of GC patients with different AJCC tumor stages (I-II v.s. III-IV). (C) The receiver operating characteristic (ROC) curves for predicting patient survival using AJCC stage (left panel), OCT1 level (middle panel) or combination of two factors (right panel). The area under curve (AUC) and the corresponding 95% CI are shown in the plots. (D) Different factors (including OCT1, AJCC stage, tumor size, sex and age) were analyzed for their association with patient survival using Cox regression model. The hazard ratio and 95% confident interval (CI) are plotted for each factor. (E) The GC patients in the Singapore cohort were stratified according to OCT1 mRNA level (median split) and compared for the survival using Kaplan-Meier analysis. (F, G) Gain in OCT1 gene copy number associated with poor survival in both the Singapore cohort (F) and VUMC cohort (G).

performed FISH analysis on six GC tissues and found two samples with gained copy number of OCT1 gene (figure 1F). Combined analysis on the mRNA and CNV data in the TCGA cohort revealed that gained CNV of OCT1 associated with significantly higher mRNA level in GC (figure 1G). In addition, amplifications of OCT1 displayed mutual exclusivity with those of KRAS and FGFR2 genes in the RTK pathway, which are also frequently altered in GC (figure 1H).

OCT1 gene amplification and upregulation associate with poor survival of GC

We further studied the association between OCT1 expression level and different clinicopathological features of GC. When the patients were stratified according to OCT1 expression levels (median split), significantly larger tumour size was found in the OCT1-high group ($p < 0.0001$, see online supplementary figure

S2A). However, OCT1 was not associated with the sex, age, American Joint Committee on Cancer (AJCC) staging, or any histological subtypes of patients with GC (see online supplementary figure S2B, dataset shown in online supplementary table S1). Notably, OCT1 overexpression strongly associated with poor survival of patients with GC ($p < 0.0001$, Mantel-Cox test, figure 2A), and the 5-year survival rate in the OCT1-high group (8.9%) was substantially lower than that of the OCT1-low group (51.1%). In fact, the stratification by OCT1 level displayed even higher prognostic significance than the widely employed AJCC staging (significance $p = 0.0002$, 5-year survival: 20 vs 42.2 months, figure 2B). When the GC cases were stratified into four groups according to either OCT1 expression (quartile split) or AJCC staging (I-IV), a better discrimination was also achieved by the OCT1-based method (see online supplementary figure S2C,D). We determined the

receiver operating characteristic curves for the prediction of patient survival using either AJCC stage or OCT1 level, or a combination of both (figure 2C). The area under curve (AUC) of OCT1-based prediction (0.751) was higher than that of the AJCC stage-based model (0.714), while the combination of both factors achieved the highest AUC value (0.800).

Multivariate Cox regression survival analysis adjusting for AJCC stage, tumour size, sex and age of patients consistently reported strong correlation between OCT1 overexpression and shorter survival ($p=0.0001$, HR=3.27, 95% CI 1.80 to 5.92, figure 2D; also see online supplementary figure S2E). We validated the prognostic significance using the Singapore cohort and found downregulation of OCT1 mRNA significantly associated with survival in patients with GC ($p=0.007$, Kaplan–Meier analysis, figure 2E).²¹ In addition, the gained CNV of OCT1 was also significantly associated with shorter survival, as revealed by the Singapore cohort ($p=0.015$, figure 2F) and the VUMC GC cohort ($p=0.040$, figure 2G).²⁰ These findings consistently suggest OCT1 as a promising prognostic marker for GC cells.

Identification of ERK signalling pathway as regulatory target of OCT1

As described above, the exact pathways that OCT1 may regulate in cancers remain unclear. To probe the OCT1-associated pathways on an unbiased basis, we performed GSEA using high throughput RNA-sequencing data of the GC cohort of The Cancer Genomic Atlas project (TCGA, 282 patients) and a multi-tumour dataset (including 64 human tumours and 27 normal tissue samples) from the Gene Expression Omnibus database.²² GSEA is designed to detect coordinated differences in expression of predefined sets of functionally related genes.²³ Among all the 189 predefined ‘oncogenic signature’ gene sets, the PDGF/ERK pathway was identified with the strongest association with OCT1 expression in the TCGA GC dataset (figure 3A) and the multi-tumour dataset (figure 3B). In addition, the Enrichment Map algorithm adjusting for gene set redundancy also identified the RAS-ERK signalling pathway in strong correlation with OCT1 expression (see online supplementary figure S3). Of note, approximately half (2083/4348) of OCT1-correlated genes were found as inducible targets of ELK1,²⁴ which is a major effector in the ERK signalling pathway (figure 2C). These findings consistently suggest that OCT1 may be involved in the activation of the ERK signalling pathway.

OCT1 expression associates with ERK phosphorylation in GC

In light of the association between OCT1 and ERK signalling, we explored whether OCT1 may correlate with ERK2 expression level or its phosphorylation/activation in GC tissues. Interestingly, analysis on RNA-sequencing data suggested no association between OCT1 and ERK2 mRNA levels (figure 3D), while the tissue immunofluorescence study revealed a strong correlation between OCT1 expression and the level of ERK2 phosphorylation (pERK2) in GC (figure 3E). In addition, the levels of OCT1 and pERK2 were coordinately increased from normal tissues to GC tissues (figure 3F,G). These results suggest potential implication of OCT1 in the phosphorylation/activation of ERK2 in GC.

Regulation of synbindin-mediated ERK phosphorylation by OCT1

Since our group has identified synbindin as a pivotal regulator of ERK2 phosphorylation in GC,⁹ we questioned whether OCT1 might affect ERK signalling by regulating synbindin. This

assumption was well supported by the putative OCT1-binding sites in the synbindin promoter (within –700 bp upstream of the transcription starting site (TSS); see online supplementary figure S4A). Luciferase reporter assay suggested this promoter region could be transactivated by OCT1 (see online supplementary figure S4B). Further, we performed ChIP assay to test the binding of OCT1 to synbindin promoter in vivo. As a result, the promoter region of synbindin was amplifiable from the DNA recovered from the immunoprecipitation complex using a specific antibody for OCT1 but not a control IgG (see online supplementary figure S4C), thus confirming the binding of OCT1 on the promoter regions of synbindin. To identify the exact binding site of OCT1 on synbindin promoter, we developed a sequence scanning approach using the 3DTF method²⁵ based on the crystallographic data on the OCT1 protein–DNA complex (pdb entry: 1O4X). The sequence ranging from –253 to –241 upstream of TSS of synbindin was found with the highest binding propensity to OCT1 (figure 4A). To confirm this finding, we analysed the enrichment of OCT1-binding stretches derived from previous ChIP-seq study²⁶ in the synbindin promoter. This method also identified the –253 to –241 region as the strongest binder to OCT1 (figure 4B). Finally, a luciferase reporter inserted by this short sequence could be strongly transactivated by OCT1 (figure 4C), while other sequences mapping to non-predicted sites were not transactivated (see online supplementary figure S4D,E). These results confirmed the findings of the two dedicated prediction methods, indicating that OCT1 binds to and transactivates –253 to –241 region of the synbindin promoter.

In support of the effect of OCT1 on transactivating synbindin promoter, ectopic expression of OCT1 markedly upregulated synbindin expression at mRNA and protein levels (figure 4D,E). Consistently, inhibition of OCT1 by specific siRNAs significantly decreased mRNA and protein levels of synbindin (figure 4F,G). The expression levels of OCT1 and synbindin showed similar trends in different GC cell lines, which supported the regulatory effect of OCT1 on synbindin (figure 4H,I). Accordingly, the level of phosphorylated ERK (p-ERK) was also higher in the OCT1 and synbindin-expressing cells (figure 4H), although the genetic background of AGS (KRAS mutation) and MKN45 (MET amplification) should also be taken into account. Ectopic expression of OCT1 significantly increased synbindin level and ERK2 phosphorylation in GC cells (figure 4J). The subtle change on total ERK2 level confirmed that OCT1 has little effect on ERK2 expression. Taken together, these results indicate that OCT1 regulates ERK2 phosphorylation/activation by transactivating synbindin in GC cells.

OCT1 as a determinant of synbindin expression in GC tissues

Further, we performed immunofluorescence assay to analyse the expression levels of OCT1 and synbindin in normal gastric mucosa and GC tissues. As shown in figure 5A–C, OCT1 and synbindin protein levels showed remarkable correlation ($p<0.0001$, Pearson correlation). Consistently, RNA-sequencing data revealed significant correlation between the mRNA levels of OCT1 and synbindin across different cancer types and normal tissues ($p<0.0001$, Pearson correlation, figure 5D). The relationship between OCT1 and synbindin mRNAs in GC tissues was further analysed using microarray data,²⁷ which suggested strong correlation between mRNA levels of the two genes ($p=0.0002$, Pearson correlation, figure 5E). These results, based on human cancer tissues, strongly suggest that OCT1 is a determinant of synbindin expression in GC cells.




Figure 3 Identification of extracellular signal-regulated kinase (ERK) signalling as a regulatory target of octamer transcription factor 1 (OCT1). (A, B) Gene Set Enrichment Analysis (GSEA) identified significant association between OCT1 and ERK signaling pathway in both the TCGA gastric cancer dataset and the multi-tumor dataset. (C) Circos plot showing 2083 of the 4348 OCT1-associated genes are inducible by ELK1, a major effector gene in ERK signaling pathway. The histogram shows corresponding P values (log1/2 transformed) of the associated genes, with red color representing ELK1-inducible genes while blue indicating non-inducible genes. (D) The expression levels of OCT1 and ERK2 were subjected to Pearson correlation analysis, which suggested no correlation between the two genes (P=0.97). (E) Correlation between the protein levels of OCT1 and phosphorylated ERK (pERK) as determined by immunostain of gastric cancer and normal tissues. (F) Representative immunofluorescent image of OCT1 (labeled in red) and pERK (green) in normal gastric tissues. Zoomed sections are shown on the right panels. Scale bars indicate 100µm in all panels. (G) Representative gastric cancer immunostain images acquired by the same procedures as described previously.

Synbindin-dependent effects of OCT1 in GC cells

To test whether the effects of OCT1 on ERK phosphorylation/activation is dependent on synbindin expression, the human GC MGC803 cells were transfected with OCT1 expression vector

or control vector, in the absence or presence of siRNAs specific for synbindin. While OCT1 significantly increased the levels of pERK and pELK1 (without affecting p-MEK level), knockdown of synbindin efficiently suppressed the effects of OCT1 (figure 5F).




Figure 4 Transactivation of synbindin by octamer transcription factor 1 (OCT1) and effects on extracellular signal-regulated kinase (ERK) activation. (A) TF-DNA crystallography structure-based prediction of OCT1 binding sites on synbindin promoter (OCT1-DNA complex structure shown on the right). Arrow heads in red and in grey respectively indicate the predicted binding sequence and other sequences tested by luciferase assay. (B) ChIP-seq enriched motif-based prediction of OCT1 binding on synbindin promoter (analysis scheme shown in the right panel). (C) The predicted OCT1-binding sequence (-253 to -241) and its mutated variant (both sequences indicated) were respectively inserted into a luciferase reporter and analyzed for their responses to OCT1. (D) Gastric cancer cells were transfected with OCT1 or control vector, and the mRNA level of synbindin was quantified by qPCR. (E) Western Blot of OCT1 and synbindin of cells treated as described as above. (F) Cells were treated by OCT1 siRNAs and control siRNA, and the level of synbindin mRNA was quantified by qPCR. The 18S rRNA level was used for normalization. (G) Western Blot showing decrease of synbindin protein upon knockdown of OCT1 using specific siRNAs. (H) Western blot showing the expression levels of OCT1, synbindin, p-ERK and ERK in five gastric cell lines. OCT1 expression is upregulated in GC cells as compared to normal gastric mucosa cell GES-1. (I) The mRNA level of synbindin in gastric cell lines was detected by RT-qPCR. (J) Cells were transfected by OCT1 or control vector, followed by Western Blot analysis using specific antibodies for synbindin, phosphorylated ERK (pERK) or total ERK.

These results supported our notion that synbindin plays a central role in mediating OCT1-induced ERK signalling.

Moreover, we found that synbindin is required for the effects of OCT1 on the malignant behaviours of GC cells. Transwell assay revealed a significant increase in the invasiveness of OCT1-expressing cells, which could be suppressed by knockdown of synbindin (figure 5F,G). In addition, MTT assay suggested pro-proliferation effect of OCT1 on GC cells (figure 5H), and flow cytometry combined with annexin V-FITC staining indicated the strong anti-apoptotic effect of

OCT1 (figure 5I,J). Importantly, suppression of synbindin expression by specific siRNAs impaired these pro-malignant effects of OCT1 (figure 5F-J). Since the MKN45 cells expressed a higher level of OCT1, we examined the effect of suppressing OCT1 on cell proliferation and invasion. As a result, knockdown of OCT1 using specific siRNAs inhibited the proliferation and invasion of MKN45 cells, while ectopic expression of synbindin restored cell proliferation and invasion (figure 5K,L). These results suggest that OCT1 contributes to malignant behaviours of GC cells by regulating synbindin expression.




Figure 5 Roles of octamer transcription factor 1 (OCT1) in regulating synbindin expression and cell aggressiveness. (A) Immunofluorescent detection of OCT1 and synbindin expression in normal gastric tissues. Scale bars indicate 20 μ m in all panels. (B) Immunostain of OCT1 and synbindin in human gastric cancer tissues. (C) Correlation between the protein levels of OCT1 and synbindin in human gastric cancer tissues as determined by immunofluorescence. (D) The mRNA levels of OCT1 and synbindin determined by RNA-sequencing (GSE28866) showed strong correlation in cancer and normal tissues. (E) The OCT1 and synbindin mRNA levels detected by microarray (GSE27342) were strongly correlated in human gastric tissues. (F) Representative Transwell assays of MGC803 cells stably expressing the control vector or OCT1 vector treated without/with the specific siRNAs for synbindin. Scale bars indicate 100 μ m in all panels. (G) Statistical result based on three independent Transwell assays. (H) Cell Counting Kit-8 (CCK8) assay of MGC803 cells stably expressing control vector or OCT1 vector without/with treatment by synbindin siRNAs. (I) Apoptosis of cells treated by above-mentioned procedures as measured by flow cytometry combined with phycoerythrin (PE)-conjugated annexin V staining. (J) Statistics based on 3 independent apoptotic assays (** $P < 0.01$; *** $P < 0.001$, t-test). (K) The MKN45 cells were treated by two specific siRNAs (left and right panels) without or with ectopic expression of synbindin, and cell proliferation was measured by CCK8 assay (OD450 absorbance). (L) Effects of OCT1 silencing and synbindin ectopic expression on the invasiveness of MKN45 cells. Scale bars indicate 100 μ m in all panels.

OCT1-synbindin signalling promotes ELK1 and RSK phosphorylation

Previous studies suggested that ERK signalling can be spatially separated into substrates that bind to the DEF (docking site for ERK, FXFP) domain or the D (docking) domain.²⁸ To determine the specific ERK substrates that are affected by OCT1 and synbindin, we used Western blot to detect the phosphorylation of ELK1 (DEF-domain substrate) and RSK (D-domain substrate) in MGC803 cells expressing ectopic OCT1. Interestingly, we found

that the phosphorylation of ELK1 and RSK were both increased upon OCT1 upregulation, but the change of p-ELK1 was more pronounced (figure 6A). In support of this, OCT1 silencing in MKN45 cells substantially decreased ELK1 phosphorylation, while synbindin ectopic expression largely restored the p-ELK1 level. In contrast, OCT1 silencing only induced a mild decrease of p-RSK level (see online supplementary figure S5).

Accordingly, knockdown of ELK1 also abrogated the effect of OCT1 overexpression on cell proliferation, but RSK silencing




Figure 6 Effects of octamer transcription factor 1 (OCT1) on the phosphorylation of extracellular signal-regulated kinase (ERK) and substrate proteins. (A) Cells were transfected with OCT1 expression vector in the absence or presence of siRNAs specific for synbindin (si-synbindin #1 & #2), and the levels of pERK/ERK, p-ELK1/ELK1, p-RSK/RSK were detected by Western Blot. The α -tubulin level was used as loading control. (B) Effects of ectopic OCT1 expression and knockdown of ERK substrates (RSK and ELK1) on cell proliferation at the last time point (48h) using t-test (* $P < 0.05$, ** $P < 0.01$). (C) The Hex spherical polar Fourier protein docking algorithm was employed to probe the synbindin-ERK complex conformation based on the crystallography structures of synbindin (PDB ID code 3EAP) and ERK2 (PDB ID code 2J0Z). (D) Input amount of the ERK mutants in D-domain (mut-D, D319N) and in DEF domain (mut-DEF, L198A, L232A, and Y261A) for GST pull-down assay. (E) The co-immunoprecipitated synbindin protein under each condition is shown in the Western Blot.

only mildly affected the effect of OCT1 (figure 6B). Taken together, our data suggest that ELK1 (docked to DEF domain) plays more important roles in mediating the pro-malignant potentials of OCT1/synbindin.

Mode of synbindin-ERK interaction

We found previously that synbindin binds to MEK and ERK on the Golgi, facilitating ERK phosphorylation by MEK.⁹ In addition, the C-terminal Longin (LDc) domain of synbindin was found to interact with ERK protein. Here we further probe the structural motif of ERK that interacts with synbindin by protein-protein docking combined with experimental validation. The Hex spherical polar Fourier protein docking algorithm²⁹ was implemented to determine the complex conformation of synbindin (PDB ID code 3EAP) and ERK2 (PDB ID code 2J0Z). Intriguingly, the synbindin LDc domain perfectly matched with the ERK DEF domain in terms of geometry and electrostatic complementation (figure 6C). The binding

mode resembled the interaction between ERK and PEA-15 (an ERK spatial regulator)³⁰ and a designed ankyrin repeat protein³¹ that has recently been determined by crystallography (see online supplementary figure S6A–C). Since the ERK amino acid residues Leu198, Leu232 and Tyr261 were found to be important for DEF docking,²⁸ we proposed they might also be involved in the binding to synbindin. To test this, we fused ERK to glutathione S-transferase (GST) and generated mutations to disrupt the above residues (L198A, L232A and Y261A). The GST pull-down assay identified interaction between synbindin and wild-type ERK, which was abolished by mutations in the DEF domain (figure 6D,E). However, the control mutation D319N sitting in the D-domain did not affect the interaction between synbindin and ERK. These data suggest that synbindin binds to the DEF domain of ERK rather than the D domain. Protein docking suggested that synbindin-ERK interaction facilitates subsequent docking of MEK1 to the ERK activation loop (predicted model shown in online supplementary figure S6D),




Figure 7 Octamer transcription factor 1 (OCT1) promotes synbindin expression and extracellular signal-regulated kinase (ERK) phosphorylation in vivo. (A) The right panel shows statistical analysis of OCT1 expression based on multiple xenograft tumors ($P < 0.01$, two-sided student t-test). (B) Representative image and statistical analysis of synbindin expression in xenograft tumors expressing OCT1 or control vector. Synbindin is stained using specific antibody in red. Scale bars indicate 10 μ m. (C) The level of phosphorylated ERK (pERK) was dramatically increased by the expression of OCT1. (D) Western Blot of xenograft tissues in OCT1-expressing group and control group. (E) Growth curves of xenograft tumors expressing OCT1 or control vector. (F) Nude mice carrying xenograft tumors that expressed OCT1 or control vector. The ranges of tumors are marked by dashed lines. (G) Image of excised tumors expressing OCT1 or control vector. (H) Schematic representation for the OCT1-synbindin-ERK regulatory pathway proposed in this study. OCT1 transactivates synbindin, which is a core subunit of a tether complex on the Golgi involved in vesicle transportation. Synbindin functions as a molecular scaffold that binds to both MEK and ERK2, facilitating ERK phosphorylation on the Golgi apparatus. This promotes the activation of ERK substrates (especially the DEF-domain substrate ELK1), leading to potentiation of cell proliferation and invasion in gastric cancers.

which seems to be relevant to the enhanced activation of ERK observed in our previous study.⁹

Further, we questioned whether potential phosphorylation of synbindin may be involved in this process. By analysing the PhosphoSite database³² that integrates published phosphorylation data using unbiased high-throughput proteomic

approaches, we summarised the frequency of all detected protein phosphorylation in 32 reported GC cell lines (data in online supplementary table S2). The ERK2 protein was found as the most frequently phosphorylated protein in these GC cell lines (see online supplementary figure S6E), which is consistent with a previous report about the high prevalence of

ERK activation in GC.⁵ Although 6497 phosphorylation sites were detected in the proteome, no phosphorylation of synbindin was identified in any cell line. Thus, the phosphorylation of synbindin does not seem to be required for ERK activation in GC cells. Taken together, these findings support our notion that synbindin facilitates MEK-ERK interaction and ERK phosphorylation, leading to activation of ERK substrates (proposed model shown in online supplementary figure S6F).

Effects of OCT1 on ERK activation and tumour growth in xenograft mouse model

Finally, we tested whether upregulation of OCT1 could promote tumourigenicity of GC cells *in vivo* using a xenograft model in nude mice. Stable cell lines carrying either OCT1 expression vector or the control vector were injected subcutaneously to the right flank of nude mice to allow xenograft tumour formation. At 14 days after first injection, mice in the OCT1 group and control group were sacrificed and the xenograft tissues were subjected to Western blot and immunofluorescence analyses. As a result, OCT1 was efficiently upregulated in the OCT1 group compared with the control group (figure 7A). Meanwhile, the expression levels of synbindin (figure 7B) and phosphorylated ERK2 (figure 7C) were significantly higher in the OCT1-upregulated group. These results were also confirmed by Western blot, which revealed a substantial increase in the levels of synbindin and phosphorylated (but not total) ERK2 in the tissues of the OCT1-upregulated group (figure 7D). As a result of enhanced ERK signalling, xenograft tumour growth in the OCT1-upregulated group was significantly faster than that of the control group (figure 7E–G). These results strongly supported our notion that OCT1 promotes gastric carcinogenesis and this effect involves synbindin-mediated ERK signalling activation.

DISCUSSION

The OCT1 transcription factor, a homologue of the OCT4 pluripotency factor, was found to express in GC and its precancerous lesion. However, the functional implication and prognostic value of OCT1 in GC have been poorly defined. In this study we report recurrent amplification and upregulation of the OCT1 gene associated with poor prognosis of patients with GC. In addition, we reveal the underlying mechanism for such an association by demonstrating OCT1 as a determinant of synbindin-mediated ERK signalling and related aggressive phenotypes in GC cells.

First, our study identifies OCT1 as a promising biomarker for the diagnosis and prognosis of GC. OCT1 is significantly upregulated in GC and its precancerous lesion, and the expression of OCT1 can be conveniently detected by histochemistry of tissue biopsy obtained by gastric endoscopy. Thus, OCT1 should be further evaluated for its diagnostic value in early GC. Importantly, the strong association between OCT1 upregulation and poor outcome of patients with GC has been confirmed by our data and an independent dataset. In addition, the prognostic significance of OCT1 CNV is also supported by two independent datasets. These interesting results suggest that OCT1 overexpression (detectable by immunohistochemistry) or gained CNV (by FISH) can be used as prognostic markers in GC. The GSEA indicated that OCT1 overexpression marks activation of the RAS/ERK signalling pathway, thus may provide useful information for targeted therapy.

The recurrent amplification of the OCT1 gene and its prominent ability to increase tumour aggressiveness suggest that

OCT1 may play a driving role in gastric carcinogenesis. Thus, targeting OCT1 may present a favourable therapeutic strategy for GC cells, especially for the cases with deregulated ERK signalling. Interestingly, OCT1 has recently been reported to be associated with ERK signalling in oesophageal carcinogenesis,³³ suggesting that OCT1 may affect ERK signalling and cancer aggressiveness in a broader scenario. Thus, it warrants further studies to clarify the clinical and biological relevance of OCT1 in other types of cancers.

By demonstrating the effect of OCT1 on transactivating synbindin, we are able to provide a more complete model for the spatial regulation of ERK signalling on the Golgi apparatus (schematic representation in figure 7H). Apart from the canonical Ras-Raf-MEK-ERK signal transduction flow, recent studies have stressed the importance of ERK signalling regulation in specific subcellular compartments, a so-called 'spatial regulatory' mechanism.³⁴ In our recent studies,⁹ we identified synbindin as a key molecular scaffold that confers spatial regulation of ERK signalling on the Golgi (illustrated in figure 7H). In this study, we further identify OCT1 as an upstream regulator of synbindin in GC and provide structural insight into synbindin-mediated ERK activation. We show that synbindin LDC domain binds to the DEF domain of ERK, leading to phosphorylation of ERK substrates (especially DEF-domain substrate) and increased cell aggressiveness. Importantly, our study on GC tissue samples reported a strong correlation between OCT1 expression and ERK2 phosphorylation levels ($p < 0.0001$), suggesting that OCT1 is a crucial regulator for ERK2 activation in GC. The recurrent amplification of the OCT1 gene, together with its regulation on synbindin-related ERK signalling, suggests OCT1 as a potential driver of malignant behaviours of GC cells.

The effects of OCT1 on ERK signalling may also help to explain other reported activities of OCT1. As a homologue of OCT4 pluripotency factor, OCT1 has been found without the ability to induce pluripotency.³⁵ Instead, OCT1 was suggested to associate with the self-renewal of cancer stem cells³⁶ and the epithelial-mesenchymal transition (EMT) process.¹⁶ Interestingly, ERK signalling has also been found to be crucial for the stem-like cells in multiple cancers, including GC,³⁷ colorectal cancer,³⁸ prostate cancer,³⁹ breast cancer⁴⁰ and glioblastoma.⁴¹ Recent studies have also clearly demonstrated the important roles of ERK signalling in the EMT process.^{42–44} Thus, it warrants further study whether OCT1 may contribute to the stem-like behaviours of cancer cells and the EMT process by regulating of the ERK signalling pathway.

In conclusion, our results identify OCT1 as a determinant of synbindin-mediated activation of ERK signalling in GC, and OCT1 is a potential prognostic marker and therapeutic target in GC.

Contributors JQ and JX wrote the paper. JQ, XK, JW, HC and YH conducted the experiments. ND and PT contributed data. JX, HC, WZ and J-YF analysed data. J-YF and JX supervised the research. JQ and XK contributed equally.

Funding This project was supported by grants from the National Basic Research Program of China 973 Program (2010CB5293), the National High Technology Research and Development Program of China 863 Program (2012AA02A504), the Program for Innovative Research Team of Shanghai Municipal Education Commission, and National Natural Science Foundation of China (30971330, 31371420, 81320108024) to JY Fang. The project was also partially supported by National Natural Science Foundation of China (81000861, 81322036, and 81272383), Shanghai 'Oriental Scholars' project (2013XJ), Shanghai Science and Technology Commission 'Pujiang Project' (13PJ1405900), and Shanghai Natural Science Foundation (12ZR1417900) to J Xu. The sponsors of this study had no role in the collection of the data, the analysis and interpretation of the data, the decision to submit the manuscript for publication, or the writing of the manuscript.

Competing interests None.

Patient consent Obtained.

Ethics approval Ethics approval was provided by the Ethics Committee of the Shanghai Jiao-Tong University School of Medicine, Renji Hospital.

Provenance and peer review Not commissioned; externally peer reviewed.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 3.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/3.0/>

REFERENCES

- Coupland VH, Lagergren J, Luchtenborg M, *et al.* Hospital volume, proportion resected and mortality from oesophageal and gastric cancer: a population-based study in England, 2004–2008. *Gut* 2013;62:961–6.
- Camargo MC, Kim WH, Chiaravalli AM, *et al.* Improved survival of gastric cancer with tumour Epstein-Barr virus positivity: an international pooled analysis. *Gut* 2014;63:236–43.
- Shibata W, Ariyama H, Westphalen CB, *et al.* Stromal cell-derived factor-1 overexpression induces gastric dysplasia through expansion of stromal myofibroblasts and epithelial progenitors. *Gut* 2013;62:192–200.
- Wang S, Wu X, Zhang J, *et al.* CHIP functions as a novel suppressor of tumour angiogenesis with prognostic significance in human gastric cancer. *Gut* 2013;62:496–508.
- Paterson AL, Shannon NB, Lao-Sirieix P, *et al.* A systematic approach to therapeutic target selection in oesophago-gastric cancer. *Gut* 2013;62:1415–24.
- Deng N, Goh LK, Wang H, *et al.* A comprehensive survey of genomic alterations in gastric cancer reveals systematic patterns of molecular exclusivity and co-occurrence among distinct therapeutic targets. *Gut* 2012;61:673–84.
- Dulak AM, Schumacher SE, van Lieshout J, *et al.* Gastrointestinal adenocarcinomas of the esophagus, stomach, and colon exhibit distinct patterns of genome instability and oncogenesis. *Cancer Res* 2012;72:4383–93.
- Kholodenko BN, Hancock JF, Kolch W. Signalling ballet in space and time. *Nat Rev Mol Cell Biol* 2010;11:414–26.
- Kong X, Qian J, Chen LS, *et al.* Synbindin in extracellular signal-regulated protein kinase spatial regulation and gastric cancer aggressiveness. *J Natl Cancer Inst* 2013;105:1738–49.
- Lee MC, Toh LL, Yaw LP, *et al.* Drosophila octamer elements and Pdm-1 dictate the coordinated transcription of core histone genes. *J Biol Chem* 2010;285:9041–53.
- Wang VE, Tantin D, Chen J, *et al.* B cell development and immunoglobulin transcription in Oct-1-deficient mice. *Proc Natl Acad Sci U S A* 2004;101:2005–10.
- Bertolino E, Reddy K, Medina KL, *et al.* Regulation of interleukin 7-dependent immunoglobulin heavy-chain variable gene rearrangements by transcription factor STAT5. *Nat Immunol* 2005;6:836–43.
- Shakya A, Cooksey R, Cox JE, *et al.* Oct1 loss of function induces a coordinate metabolic shift that opposes tumorigenicity. *Nat Cell Biol* 2009;11:320–7.
- Jin T, Li H. Pou homeodomain protein OCT1 is implicated in the expression of the caudal-related homeobox gene Cdx-2. *J Biol Chem* 2001;276:14752–8.
- Almeida R, Almeida J, Shoshkes M, *et al.* OCT-1 is over-expressed in intestinal metaplasia and intestinal gastric carcinomas and binds to, but does not transactivate, CDX2 in gastric cells. *J Pathol* 2005;207:396–401.
- Hwang-Verslues WW, Chang PH, Jeng YM, *et al.* Loss of corepressor PER2 under hypoxia up-regulates OCT1-mediated EMT gene expression and enhances tumor malignancy. *Proc Natl Acad Sci U S A* 2013;110:12331–6.
- Ethell IM, Hagihara K, Miura Y, *et al.* Synbindin, a novel syndecan-2-binding protein in neuronal dendritic spines. *J Cell Biol* 2000;151:53–68.
- Summersgill B, Clark J, Shipley J. Fluorescence and chromogenic in situ hybridization to detect genetic aberrations in formalin-fixed paraffin embedded material, including tissue microarrays. *Nat Protoc* 2008;3:220–34.
- Goldman M, Craft B, Swatloski T, *et al.* The UICC cancer genomics browser: update 2013. *Nucleic Acids Res* 2013;41:D949–54.
- Buffart TE, Carvalho B, van Grieken NC, *et al.* Losses of chromosome 5q and 14q are associated with favorable clinical outcome of patients with gastric cancer. *Oncologist* 2012;17:653–62.
- Ooi CH, Ivanova T, Wu J, *et al.* Oncogenic pathway combinations predict clinical prognosis in gastric cancer. *PLoS Genet* 2009;5:e1000676.
- Brunner AL, Beck AH, Edris B, *et al.* Transcriptional profiling of long non-coding RNAs and novel transcribed regions across a diverse panel of archived human cancers. *Genome Biol* 2012;13:R75.
- Subramanian A, Kuehn H, Gould J, *et al.* GSEA-P: a desktop application for Gene Set Enrichment Analysis. *Bioinformatics* 2007;23:3251–3.
- Odrowaz Z, Sharrocks AD. ELK1 uses different DNA binding modes to regulate functionally distinct classes of target genes. *PLoS Genet* 2012;8:e1002694.
- Gabbouline R, Eckweiler D, Kel A, *et al.* 3DTF: a web server for predicting transcription factor PWMs using 3D structure-based energy calculations. *Nucleic Acids Res* 2012;40:W180–5.
- Ferraris L, Stewart AP, Kang J, *et al.* Combinatorial binding of transcription factors in the pluripotency control regions of the genome. *Genome Res* 2011;21:1055–64.
- Cui J, Chen Y, Chou WC, *et al.* An integrated transcriptomic and computational analysis for biomarker identification in gastric cancer. *Nucleic Acids Res* 2011;39:1197–207.
- Dimitri CA, Dowdle W, MacKeigan JP, *et al.* Spatially separate docking sites on ERK2 regulate distinct signaling events in vivo. *Curr Biol* 2005;15:1319–24.
- Ritchie DW, Venkatraman V. Ultra-fast FFT protein docking on graphics processors. *Bioinformatics* 2010;26:2398–405.
- Mace PD, Wallez Y, Egger MF, *et al.* Structure of ERK2 bound to PEA-15 reveals a mechanism for rapid release of activated MAPK. *Nat Commun* 2013;4:1681.
- Kummer L, Parizek P, Rube P, *et al.* Structural and functional analysis of phosphorylation-specific binders of the kinase ERK from designed ankyrin repeat protein libraries. *Proc Natl Acad Sci USA* 2012;109:E2248–57.
- Hornbeck PV, Chabra I, Kornhauser JM, *et al.* PhosphoSite: a bioinformatics resource dedicated to physiological protein phosphorylation. *Proteomics* 2004;4:1551–61.
- Wang Z, Zhu S, Shen M, *et al.* STAT3 is involved in esophageal carcinogenesis through regulation of Oct-1. *Carcinogenesis* 2013;34:678–88.
- Reiterer V, Fey D, Kolch W, *et al.* Pseudophosphatase STYX modulates cell-fate decisions and cell migration by spatiotemporal regulation of ERK1/2. *Proc Natl Acad Sci U S A* 2013;110:E2934–43.
- Kang J, Shakya A, Tantin D. Stem cells, stress, metabolism and cancer: a drama in two acts. *Trends Biochem Sci* 2009;34:491–9.
- Maddox J, Shakya A, South S, *et al.* Transcription factor Oct1 is a somatic and cancer stem cell determinant. *PLoS Genet* 2012;8:e1003048.
- Khurana SS, Riehl TE, Moore BD, *et al.* The hyaluronic acid receptor CD44 coordinates normal and metaplastic gastric epithelial progenitor cell proliferation. *J Biol Chem* 2013;288:16085–97.
- Wang YK, Zhu YL, Qiu FM, *et al.* Activation of Akt and MAPK pathways enhances the tumorigenicity of CD133+ primary colon cancer cells. *Carcinogenesis* 2010;31:1376–80.
- Rybak AP, Ingram AJ, Tang D. Propagation of human prostate cancer stem-like cells occurs through EGFR-mediated ERK activation. *PLoS ONE* 2013;8:e61716.
- Ahn HJ, Kim G, Park KS. E13 stimulates proliferation, drug resistance, and cancer stem cell properties of breast cancer cells via a MEK/ERK-dependent signaling pathway. *Biochem Biophys Res Commun* 2013;437:557–64.
- Nakada M, Nambu E, Furuyama N, *et al.* Integrin alpha3 is overexpressed in glioma stem-like cells and promotes invasion. *Br J Cancer* 2013;108:2516–24.
- Zhang W, Mendoza MC, Pei X, *et al.* Down-regulation of CMTM8 induces epithelial-to-mesenchymal transition-like changes via c-MET/extracellular signal-regulated kinase (ERK) signaling. *J Biol Chem* 2012;287:11850–8.
- Ha GH, Park JS, Breuer EK. TACC3 promotes epithelial-mesenchymal transition (EMT) through the activation of PI3 K/Akt and ERK signaling pathways. *Cancer Lett* 2013;332:63–73.
- Elsum IA, Martin C, Humbert PO. Scribble regulates an EMT polarity pathway through modulation of MAPK-ERK signaling to mediate junction formation. *J Cell Sci* 2013;126:3990–9.



OCT1 is a determinant of synbindin-related ERK signalling with independent prognostic significance in gastric cancer

Jin Qian, Xuan Kong, Niantao Deng, Patrick Tan, Haoyan Chen, Jilin Wang, Zhaoli Li, Ye Hu, Weiping Zou, Jie Xu and Jing-Yuan Fang

Gut 2015 64: 37-48 originally published online April 9, 2014
doi: 10.1136/gutjnl-2013-306584

Updated information and services can be found at:
<http://gut.bmj.com/content/64/1/37>

Supplementary Material

Supplementary material can be found at:
<http://gut.bmj.com/content/suppl/2014/04/09/gutjnl-2013-306584.DC1.html>

These include:

References

This article cites 44 articles, 26 of which you can access for free at:
<http://gut.bmj.com/content/64/1/37#BIBL>

Open Access

This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 3.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/3.0/>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

[Open access](#) (171)
[Pancreatic cancer](#) (618)

Notes

To request permissions go to:
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:
<http://group.bmj.com/subscribe/>